

Virginia Cooperative Extension

Knowledge for the CommonWealth



York County Office 100 County Drive PO Box 532 Yorktown, Virginia 23690-0532 (757) 890-4940 / Fax: 890-4033 E-Mail: ex199@vt.edu Web: www.yorkcounty.gov/vce

LAWN MAINTENANCE CALENDAR - PENINSULA AREA 2005-2006 Researched and Recommended Bermuda & Zovsia Grass (Warm Season Grasses) Bermuda & Zoysia

Activity

OVERSEED. Bermuda grass with annual ryegrass, if desired, at 5 lbs. per

1,000 sq. ft. Zoysia does not lend itself to over-seeding.

November 15 – WINTER WEED CONTROL. Important to control chickweed, dandelion,

December 15 henbit, buttercut, and many other annuals. Use combination of 2,4-D and Dicamba or

MCPP. Read label closely for use on Zoysia grass.

PRE-EMERGENT CRABGRASS CONTROL. March -Use Pendimethalin, Benefin (Balan), Mid-March

Dithiopyr(Dimension), or DCPA(Dacthal). Avoid Siduron (Tupersan, Trey). Make the first application when forsythia is in the first week of bloom. The second application is product

dependent; read and follow the label carefully.

February 15 -2ND ANNUAL WINTER WEED INSPECTION AND CONTROL, if

March 15 necessary. 2, 4-D and Dicamba herbicide will control most wide leaf weeds.

April 1 <u>Fertilize</u> – Obtain a soil test every 3 or 4 years and use the recommendations in the most

recent report to determine the N-P-K rates for your lawn (e.g., 25-3-7 or 12-4-8).

Apply a complete fertilizer in sufficient amount to provide 1 ½ lbs. of actual N (nitrogen).

Example: 15 lbs. of 10-10-10 or 18 lbs. of 8-8-8 per 1,000 sq. ft. or

the equivalent of another analysis.

See other programs in Extension Publication 452-238 (Soil Test

Notes) or 430-011 (Lawn Fertilization in Virginia).

2005-2006 Recommended Bermudagrass Varieties

(Varieties with the notation # are only recommended in Southeastern Virginia)

Midfield, Midiron, Midlawn, Patriot, Quickstand, Shanghai[#], Tifgreen[#], TifSport[#], Tifway II[#], Tufcote and Vamont.

2005-2006 Recommended Zoysiagrass Varieties

Meyer (vegetatively propogated) Zenith (seeded)

May 1 – June 1

OVERSEEDING. 1-2 lbs. of hulled Bermuda on common Bermuda lawns if needed (deadline-July 1). Soil preparation via thatching or raking is necessary for seed establishment.

FERTILIZE. Apply equivalent of 1 to 1 ½ lbs. of actual nitrogen per 1,000 sq. ft.

EXAMPLE: 10 to 15 lbs. of 10-10-10 or 12 to 18 lbs. of 8-8-8 per 1000 sq. ft. or the equivalent of other fertilizers. Table 1 in publication 430-011 has other analyses listed.

May 1 – June 1

WATERING. Begin deep watering every 7-10 days, if necessary.

CUTTING HEIGHT. Approximately 1" throughout season.

THATCH. May 20 – June 15. (if needed)

July 1

FERTILIZE. Apply equivalent of 1 to 1 ½ lbs. of actual nitrogen per 1,000 sq. ft. Omit on Zoysia.

July 1 – August 1 **WEED CONTROL.** Apply a combination of 2,4-D and Dicamba or MCPP herbicide to control annual summer wide leaf weeds, if needed.

WATERING. Continue to deep water every 7-10 days, if weather necessitates. Avoid night watering – turf should "dry off" before night.

DISEASE AND INSECT ALERT. Diseases and insects are few with these species of grasses, but because of temperature and humidity, a disease could cause problems. Contact the Extension Office for problems in turf.

Other helpful publications available from the Extension Office or at the website

www.ext.vt.edu/resources/

Publication No.	426-609	Selecting Landscape Plants – Ground Covers
	430-009	Lawn Establishment In Virginia
	427-045	Lawn Weed Control
	430-011	Lawn Fertilization in Virginia
	425-125	Soil Test Box Kit (Extension Office ONLY)